

DYNAMICS WITH A NON-STANDARD INERTIA-ACCELERATION RELATION: AN ALTERNATIVE TO DARK MATTER IN GALACTIC SYSTEMS

Mordehai Milgrom

Department of Physics
Weizmann Institute of Science
76100 Rehovot, Israel

ABSTRACT We begin to investigate particle dynamics that is governed by a nonstandard kinetic action of a special form. We are guided by a phenomenological scheme—the modified dynamics (MOND)—that imputes the mass discrepancy, observed in galactic systems, not to the presence of dark matter, but to a departure from Newtonian dynamics below a certain scale of accelerations, a_o . The particle’s equation of motion in a potential ϕ is derived from an action, S , of the form $S \propto S_k[\vec{r}(t), a_o] - \int \phi dt$. The limit $a_o \rightarrow 0$ corresponds to Newtonian dynamics, and there the kinetic action S_k must take the standard form. In the opposite limit, $a_o \rightarrow \infty$, we require $S_k \rightarrow 0$ —and more specifically, for circular orbits $S_k \propto a_o^{-1}$ —in order to attain the phenomenological success of MOND. If, to boot, S_k is Galilei invariant it must be time-non-local; indeed, it is non-local in the strong sense that it cannot even be a limit of a sequence of local, higher-derivative theories, with increasing order. This is a blessing, as such theories need not suffer from the illnesses that are endemic to higher-derivative theories. We comment on the possibility that such a modified law of motion is an effective theory resulting from the elimination of degrees of freedom pertaining to the universe at large (the near equality $a_o \approx cH_o$ being a trace of that connection). We derive a general virial relation for bounded trajectories. Exact solutions are obtained for circular orbits, which pertain to rotation curves of disk galaxies: The orbital speed, v , and radius, r , are related by (the rotation curve) $\mu(a/a_o)a = \frac{d\phi}{dr}$ where $a = v^2/r$, and $\mu(x)$ is simply related to the value of the kinetic action, S_k , for a circular trajectory. Expressions for the angular momentum and energy of circular orbits are derived. Beyond the obvious matter-of-principle differences between the modified-law-of-motion theories treated here, and modifications of gravity, there are important applicative differences. For example, the expression for the angular momentum—an invariant under temporal variations of a_o —is different, and the evolution of disk galaxies, expected to ensue such

variations, would be different in the two classes of theories. The various points we make are demonstrated on examples of theories of this class, but we cannot yet offer a specific theory that satisfies us on all accounts, even at the non-relativistic level that we consider here. We also explore, in passing, theories that depart from the conventional Newtonian dynamics for very low frequencies. Such theories may be written that are linear, and hence more amenable to solution and analysis than the MOND theories; but, they do not fare as well in explaining the observations.

I. INTRODUCTION: THE GALACTIC MASS DISCREPANCY AND THE MODIFIED DYNAMICS

It has been posited that the observed mass discrepancy, evinced by galactic systems, stems from the breakdown of the laws that are used to analyze their dynamics. This contrasts with the prevalent dark-matter doctrine that imputes the discrepancy to the presence of large quantities of, yet unobserved, matter. In particular, the problem has been attributed[1-3] to a departure from Newton laws that occurs in the limit of small accelerations, such as are found in galactic systems: There exists a certain acceleration parameter, a_o , much above which Newton laws apply with great accuracy. Much below a_o the dynamics (of gravitational systems, at least) departs drastically from these laws, and may be epitomized by a relation of the form

$$a^2/a_o = MG/r^2, \quad (1)$$

which gives the gravitational acceleration, a , of a test particle, at a distance r from a (point) mass M (instead of the usual $a = MG/r^2$). There are different theories that might spring from this seed relation. In fact, our purpose here is to point a new direction for searching such a theory.

Arguably, the performance of this modified nonrelativistic dynamics (MOND) has been quite successful in reproducing the observations of galactic systems [4-7] (if not altogether without dispute—see e.g. Lake[8], and the rebuttal by Milgrom[9]).

The following predictions are particularly clear-cut and well-nigh oblivious to the details of the underlying theory. Most of them pertain to circular motion in axisymmetric disk galaxies.

1. The speed on a circular orbit, supported gravitationally by a central body, becomes independent of the orbital radius for large radii[2]. This implies universally flat asymptotic rotation curves of isolated disk galaxies, in accord with high quality observations of extended rotation curves[4,5,7].
2. The asymptotic rotational speed, V_∞ , depends only on the total mass, M , of the central body via $V_\infty^4 = MGa_o$. This leads to a relation of the form $V_\infty^4 \propto L$, where L is any luminosity parameter that is proportional to the total mass. This is to be compared with the observed strong $V_\infty - L$ correlation, known as the Tully-Fisher relation[10].
3. If one approximates elliptical galaxies by isothermal spheres, MOND predicts [11] a strong correlation between the velocity dispersion in a galaxy,

σ , and its total mass M : $\sigma^4 \sim M G a_o$, reproducing the observed Faber-Jackson[12] relation between σ and the luminosity.

4. MOND predicts[2] that galaxies with very low surface density—which correspond to low mean accelerations—will evince a large mass discrepancy. This has been forcibly born out by many recent observations of low-surface-brightness disks[13], and, with less confidence, for dwarf-spheroidal satellites of our galaxy[14].

5. According to MOND, elliptical galaxies—as modeled by isothermal spheres—cannot have a mean acceleration much exceeding a_o (or a mean surface density much exceeding the critical surface density, $\Sigma_o \equiv a_o G^{-1}$). This explains the observed cutoff in the distribution of mean surface brightnesses of elliptical galaxies, known as the Fish law[15-16]

6. It is known that bare, thin disks, supported by rotation, are unstable to bar formation, and eventual breakup. MOND endows such disks with added stability that enables them to survive, provided they fall in the MOND regime, i.e. their mean acceleration is at a_o or below[17]. This can explain the observed marked paucity of galactic disks with mean surface density larger than the above Σ_o , known as the Freeman law (see ref. [16]) for a discussion of this issue).

7. Above all, detailed analysis of rotation curves, for galaxies with high-quality data, show a very close match of the predictions of MOND with the observed rotation curves of these galaxies[4,5,7].

Generally, the dynamics of systems of galaxies—such as small groups, galaxy clusters, and the Virgo infall—which are not as well understood, can also be explained by MOND, with no need to invoke dark matter[3].

In the framework of MOND, all these tell us that a modification is called for when dealing with bound orbits. There is no observational information on unbound (scattering) trajectories, for massive particles. However, we may gain insight from observations of gravitational lensing of light from distant galaxies, by foreground clusters of galaxies[18]. These however concern relativistic trajectories, which require a relativistic extension of MOND for a proper treatment (see below and Sec.IV). Henceforth we assume MOND for all types of nonrelativistic trajectories.

The value of a_o has been determined, with concurrent results, by several independent methods; these rest on the twofold role of a_o as the borderline acceleration (based on predictions 5 and 6 above), and as the scale of acceleration in the deep MOND limit (based on predictions 2 and 3). The best determination comes from the detailed studies of rotation curves[4,5,7], and is

$$a_o \sim (1 - 1.5) 10^{-8} \text{ cm s}^{-2}, \quad (2)$$

assuming the value of 75 km/s/Mpc for the Hubble constant. Values of a/a_o as small as 0.1 have been probed in studies of rotation curves; for the dynamics of the infall into the Virgo cluster one has typically $a/a_o \approx 0.01$.

MOND, which at the moment, is but a phenomenological theory, may be interpreted as either a modification of inertia, or as a modification of gravity

that leaves the law of motion intact[1]. True, General Relativity has taught us that gravity and inertia are not quite separable on the fundamental level. For instance, the Einstein field equations for the gravitational field imply the geodesic law of motion. However, at the level at which we work—i.e. a non-relativistic effective theory—it does make sense to distinguish between the interpretations. By the former we mean, foremost, a modification that must be implemented for whatever combination of forces determines the motion.

In default of a full-fledged theory Milgrom[1,2,3] had initially used a working extrapolating equation of motion of the form

$$\mu(a/a_o)\vec{a} = \vec{a}_N \equiv -\vec{\nabla}\phi, \quad (3)$$

to calculate the acceleration $\vec{a}$ from the conventional potential ϕ ($\vec{a}_N$ is the Newtonian acceleration). The extrapolating function, $\mu(x)$, satisfies $\mu(x) \xrightarrow{x \rightarrow \infty} 1$, so that Newtonian dynamics is attained in the limit of large accelerations. In the deep MOND limit we have $\mu(x) \propto x$, for $x \ll 1$ (and a_o is normalized such that $\mu(x) \approx x$), so that eq. (1) results, and all the above-mentioned predictions follow. Relation (3) is noncommittal on the matter of interpretation (see ref. [1]). As it is written it may be viewed as a modification of the law of motion for arbitrary potential ϕ , or, inverted to read $\vec{a} = -\nu(|\vec{\nabla}\phi|/a_o)\vec{\nabla}\phi$, and assumed to apply only for gravitational potential, it may be viewed, instead, as a modification of gravity.

Bekenstein and Milgrom[19] have then devised a nonrelativistic formulation of MOND that is expressly a modification of gravity: It replaces the Poisson equation ($\vec{\nabla} \cdot \vec{\nabla}\phi = 4\pi G\rho$) by

$$\vec{\nabla} \cdot [\mu(|\vec{\nabla}\phi|/a_o)\vec{\nabla}\phi] = 4\pi G\rho, \quad (4)$$

which determines the gravitational potential, ϕ , generated by the mass-density distribution ρ . The acceleration of a test particle is then given by $-\vec{\nabla}\phi$. This formulation is derivable from an action principle, enjoys the usual conservation laws, and has been shown to be satisfactory on other important accounts[19]. It reduces to eq. (3) in cases of one-dimensional symmetry, and, in fact, even for less symmetric systems such as disk galaxies, has been shown to predict very similar rotation curves as eq. (3). Notwithstanding the superior standing of eq. (4), it is eq. (3) that has been used in all the rotation-curve tests of MOND[4,5,7]—being much wieldier.

There are several relativistic theories with MOND as their non-relativistic limit[19][20]; none is without problems. They are all based on modification of gravity, in that they can be formulated as a modification of the Einstein field equations for the metric, with particles moving on geodesics. Obviously, this entails an indirect effect on other forces when gravity is important; but, inasmuch as gravity can be neglected, they do not entail a modification of the motion of particles subjected to other forces.

In this paper we begin to discuss formulations of MOND that, in contrast, may be viewed as a modification of the law of motion of a particle in a given

potential field. The main implication here is that particle dynamics is affected for whichever combination of forces is at play.

We derive the dynamics from an action principle, with an action $S = S_p + S_k$, where the potential part, S_p , takes the conventional form, while the kinetic action, S_k , carries the modification. The latter is functional of the trajectory of the particle, and is constructed using the acceleration constant a_o as the only extra parameter. The theory must approach the Newtonian dynamics in the limit $a_o \rightarrow 0$. When a_o is not much smaller than all the quantities of the dimensions of acceleration, MOND takes its effect. The deep MOND limit corresponds to $a_o \rightarrow \infty$. To obtain the behavior epitomized by eq. (1) –or, to have asymptotic flat rotation curves for isolated galaxies—we expect $S_k \propto a_o^{-1}$ in this limit, at least for circular trajectories.

Thus—in departure from the pristine statement of the MOND hypothesis—deviations from Newtonian dynamics appear, here, not only when the momentary acceleration on the trajectory is much smaller than a_o . The special role of the acceleration is highlighted only through the introduction of the constant a_o of these dimensions. On a circular, constant-speed trajectories, all the quantities with the dimensions of acceleration that can be built equal the acceleration itself. For such trajectories the acceleration only has relevancy for MOND (as is the case for rotation curves of disk galaxies).

The main purposes of this paper are: (i) To adumbrate a possible rationale for MOND, which we do in Sec. II. (ii) To demonstrate that it is possible to formulate MOND as a modification of the law of motion—in contrast to a modification of gravity—and to expound some general properties of theories based on such a modification (Sec. III). (iii) To point out significant differences between the two interpretations, and to highlight the observational implications of these differences (Sec. VI). (iv) To present general, exact results pertaining to circular motion, and point their relevance to the dynamics of disk galaxies. In particular, we give a general expression for the rotation curve of an axisymmetric disk galaxy, and derive simple expressions for the energy and angular momentum of circular orbits in Lagrangian theories; this we do in Sec. IV.

Some of the points we want to explicate become more transparent in the context of a theory where the dynamics is modified in the limit of low frequencies, instead of the limit of small accelerations. Theories of this class may be constructed that are linear, and thus more amenable to analysis and solution; alas, as potential alternatives to dark matter in galactic systems, they seem to be at some odds with the observations. We discuss these, in some detail, in Sec. V, both for their own possible merit, and as heuristic auxiliaries. Finally, we list some of the desiderata that we have not been able to implement yet, and some questions that remain open (Sec. VI).

We cannot pinpoint a specific theory. There is a large variety of theories that, as far as we have checked, are consistent with the available, relevant data (especially RCs). Also, we can offer no single model that possesses all the features we would ultimately require from a theory (see Sec. VI).

This approach opens a new course for searching for a sorely needed, relativistic extension of MOND.

II. PREAMBLE: THE POSSIBLE PROVENANCE OF THE MODIFIED DYNAMICS

Various considerations lead us to suspect that MOND is an effective approximation of a theory at a deeper stratum. Here, a most germane observation[1] is that the deduced value of a_o [eq. (2)] is of the same order as cH_o , where H_o is the present expansion rate of the universe (the Hubble constant), and c the speed of light. (For $H_o = 75 \text{ km/s/Mpc}$, $cH_o \sim 7 \times 10^{-8} \text{ cm s}^{-2}$.) This near equality may betoken an effect of the universe as a whole on local dynamics, on systems that are small on the cosmological scale.

There are, in fact, several quantities with the dimensions of an acceleration, that can be constructed from cosmological parameters[21], beside the above expansion parameter

$$a_{ex} \equiv cH_o. \quad (5)$$

Another is

$$a_{cu} \equiv c^2/R_c, \quad (6)$$

where R_c is the radius of spatial curvature of the universe; yet another is

$$a_\lambda \equiv c^2|\lambda|^{1/2}, \quad (7)$$

with λ the cosmological constant ($\lambda \equiv \Lambda/3c^2$, where Λ is sometimes used).

Only a_{ex} is observationally determined with some accuracy, as the Hubble constant has been measured to lie between about 50 and 100 km/s/Mpc (see Huchra[22] for a review). On R_c we have only a lower limit of the order of c/H_o , and on $|\lambda|$ we have only an upper limit[23-24] of order $(H_o/c)^2$. So, only upper limits exist on a_{cu} and a_λ , and these are of the order of a_{ex} . In a Friedmannian universe, with vanishing cosmological constant, one has

$$a_{cu}/a_{ex} = |1 - \Omega|^{1/2}, \quad (8)$$

where Ω is the ratio of the mean density in the universe to the closure density. (We use these relations, derived from conventional cosmology, only heuristically, as MOND may entail a modification of the cosmological equations.) So, if today Ω is not very near 1, a_{cu} and a_{ex} are comparable. If the mean mass density falls much shorter of closure—as would be in keeping with the basic premise of MOND—but a non-zero cosmological constant renders the universe flat—as would conform with inflationary models[25] (see also ref. [24])—we have today

$$a_\lambda \approx a_{ex}. \quad (9)$$

We see then that the near equality $a_o \sim a_{ex}$ may be fortuitous, with a_o being really a proxy for a_{cu} , or for a_λ (or for another cosmic parameter). The

identification of the parameter behind a_o would be an important step toward constructing an underlying theory for MOND; it is also of great impact in connection with formation of structure in the universe, and the subsequent evolution of galaxies, and galactic systems[21]: Whereas a_λ is a veritable constant, a_{ex} , and a_{cu} vary with cosmic time (in different ways). The matter has also antropic bearings.

The Newtonian limit, which corresponds to $a_o = 0$, would be realized, according to the above identifications of a_o , when the universe is static ($a_{ex} = 0$), flat ($a_{cu} = 0$), or characterized by a vanishing cosmological constant ($a_\lambda = 0$), respectively.

It is clear that to some degree the various cosmological parameters must appear in local dynamics. A finite curvature radius of the universe would enter, at least through the boundary conditions for the Einstein field equations, and will modify the local gravitational field. Likewise, a non-zero cosmological constant modifies the distance dependence of Newtonian gravity. The expansion of the universe also enters, for example by slowing down otherwise-free massive particles (or red-shifting free photons). All these are not, however, what we need for MOND. These effects are only appreciable over distances or times of cosmological scales—not relevant for non-relativistic galactic systems and phenomena. We thus do not expect H_o , R_c , and λ to enter local dynamics as a time or length scale. On the other hand, a_o is of the the same order as the galactic parameter of the same dimensions. So, if a_o does remain as a vestige of the underlying mechanism, it will be the only such parameter important for non-relativistic galactic phenomena. In contrast, *relativistic* phenomena characterized by accelerations of order a_o or less, also involve scale length of cosmological magnitude. For example, a photon trajectory passing a distance r from a mass M , such that $MG/r^2 \leq a_o$, has a curvature radius of cosmological scale (see more in Sec.IV). So, the fact that only a_o appears in MOND may be peculiar to the non-relativistic nature of this theory.

The following analogy may help illuminate the situation we have in mind: Had all our knowledge come from experiments in a small, closed laboratory on the earth’s surface, our dynamics would have involved a “universal” constant $\vec{g}$ —the free-fall acceleration. Unaware of the action of the earth, we would find a relation $\vec{F} = m(\vec{a} - \vec{g})$ between the applied force $\vec{F}$, and the acceleration $\vec{a}$ of a particle. Actually, $\vec{g}$ encapsules the effects of the mass, $M_\oplus$, and radius, $R_\oplus$, of the earth. $R_\oplus$, by itself, appears conspicuously in the dynamics only on scales comparable with itself; $\vec{g}$ alone enters the results of very-small-scale experiments characterized by accelerations not much larger then g . (The escape speed here plays the part of the speed of light in connection with MOND: $g = v_{es}^2/2R_\oplus$.) The effective departure here is of a gravitational origin, but we may describe it as a modification of inertia because it is in effect for whatever combination of forces $\vec{F}$ represents, even when there are no “local” sources of gravity (i.e. sources in addition to the earth). We would thus have no reason to consider it a modification of gravity. It may well be that the departure inherent in MOND is also of a gravitational origin (and will then automatically

preserve the weak equivalence principle in the effective theory).

But why, at all, should H_o , R_c , or λ enter local dynamics as an acceleration parameter? We can only offer a vague possible rationale: Even within the purview of conventional physics we expect a difference of sorts in the dynamics of particles with accelerations above and below a_o , if a_o stands for one of the above cosmological parameters. Consider, for example, a electric charge accelerated with acceleration a ; the lower radius of its radiation zone is $r_{rad} = c^2/a$. To say that $a \ll a_{cu}$, for example, is to say that r_{rad} is much larger than the radius of curvature of universe. In a closed universe, for example, this would mean that there is not even room for a radiation zone. Clearly, the radiation pattern must differ from the case where $a \gg a_{cu}$ for which the curvature is hardly felt within the near field. More generally, an accelerated particle carries a causal structure that includes an event horizon[26], whose scale is $\ell = c^2/a$: This is the distance to the horizon; this is the space-time distance within which a freely falling frame can be erected; this is the typical wavelength of the Unruh black-body radiation, etc.. For accelerations that are low in the context of MOND ($a \ll a_o$), the quality of the universe that is behind a_o is felt within ℓ . Conversely, we surmise that ℓ is imprinted on the conjectured inertia field (e.g. by modifying the spectrum of the vacuum fields) in a way that is then sensitive to the acceleration of a test particle interacting with this field.

All the above considerations may also usher MOND in as an effective theory of gravity.

We thus envisage inertia as resulting from the interaction of the accelerated body with some agent field, perhaps having to do with the vacuum fields, perhaps with an “inertia field” whose source is matter in the “rest of the universe”—in the spirit of Mach’s principle. If this field bears the imprint of at least one of the cosmic parameters, an effective description of dynamics in which this inertia field is eliminated, may result in a scheme like MOND.

We have not been able to put these vague ideas into a concrete use, and will have to make do, in the rest of this paper, with the study of effective theories. Nonetheless, we keep this conjectured origin before our eyes as a guide of sorts on what MOND may entail, beside the implications for the structure of galactic systems, which have been the phenomenological anchor for MOND. For example, such considerations would lead us reckon with the effects of an a_o that varies with cosmic time, as alluded to above. We may also suspect that if MOND, as we now formulate it, is only an effective theory delimited by cosmological factors, then it cannot be used to describe cosmology itself (as we would not describe the motion of earth satellites with a constant- g dynamics in the above analogy). More generally, we noted above that MOND as described above—with a_o as the only relevant parameter—may breakdown for any relativistic phenomenon with acceleration below a_o (more in Sec.VI).

There are attributes of the conventional laws of motion that we cannot take for granted in light of what we said above on the possible origin of the modified dynamics. For example, it is not obvious that the effective theory

describing MOND is derivable from an action principle. The question of which symmetries (conservation laws) the theory enjoys also becomes open. We shall certainly want to preserve the six cherished symmetries of space; to wit, the translations and rotations (or more generally the six isometries of a maximally symmetric 3-D space); what we said above raises no question with regard to these: the underpinnings of the effective theory are not expected to define a preferred position or direction. The same is true of parity. Regarding time reversal symmetry, if the expansion of the universe is in the heart of MOND, the fact that the universe has a well defined time arrow might well result in a strong TR asymmetry in the MOND regime, just as physics in an external magnetic field is TR non-symmetric. It would be very difficult to ascertain a breakdown of TR symmetry in the province of the galaxies. Many potential manifestations of this are forbidden by other symmetries, and there are other balking factors. The same is true of the time-translation symmetry. Here our guess would be that time translation might be affected through the introduction of the scale of time (H_o^{-1}), and this is not interesting, in the context of MOND, as it would lead to deviations only on very long time scales.

Similarly, we are at sea regarding full Galilei invariance. While the ultimate underlying theory may well be Galilei (Poincare) invariant, or enjoy even a higher symmetry, this is not necessarily the case for the effective theory that results from the elimination of some degrees of freedom: the surmised “inertia field”, which comprises the hidden degrees of freedom, defines a preferred frame. Perhaps the effective theory does enjoy some generalization of Galilei invariance that involves a_o as a parameter, and that reduces to Galilei invariance in the limit $a_o = 0$. Imagine, for example, an accelerated observer in a curved space. As long as its acceleration is much larger than a_{cu} of eq. (6), so that its Unruh wavelength, or the size of the region affecting the dynamics, is much smaller than the radius of curvature, he will not detect departure from Lorentz invariance due to the curvature; for $a \ll a_{cu}$ the non-zero curvature distorts the Unruh spectrum.

As phenomenology does not yet require that we relinquish any of the above symmetries, and as we do not have any concrete theoretical impetus to do so, we shall retain them in the following discussion.

We shall concentrate on effective theories of motion that are derived from an action principle, and conform to the MOND hypotheses. Such effective theories, which are obtained by eliminating some degrees of freedom from a more fundamental theory, are, in many cases, non-local; i.e., they are not derivable from an action that is an integral over a simple Lagrangian that is a function of only a finite number of time derivatives of the trajectory. A well known case in point is the Wheeler-Feynman formulation of electrodynamics[27], which results from the elimination of the electromagnetic field, and is non-local when only the particle coordinates remain as degrees of freedom. We shall show—without bringing to bear the origin of MOND— that an acceptable theory of MOND must, in fact, be strongly non-local.

III. GENERAL MOND ACTIONS

One starts by assuming that the motion of a non-relativistic test particle, in a static potential field ϕ , is governed by an equation of motion of the form

$$m\vec{A} = -\vec{\nabla}\phi, \quad (10)$$

where m is the mass of the particle, and $\vec{A}$ (replacing the acceleration $\vec{a}$ in Newtonian dynamics) depends only on the trajectory $\vec{r}(t)$ (not on the potential); its value is, in general, a functional of the whole trajectory, and a function of the momentary position on the trajectory, parameterized by the time t . Of $\vec{A}$ we further require the following: (i) That it be independent of m , so as to retain the weak equivalence principle. In what follows we take a unit mass for the particle. (ii) According to the MOND assumption, $\vec{A}$ may be constructed with the aid of the acceleration constant a_o , as the only dimensional constant. This is assumed to be the case in the non-relativistic limit. (iii) $\vec{A}$ has to approach the Newtonian expression, $\vec{a}$, in the limit $a_o \rightarrow 0$, for all trajectories, to achieve correspondence with Newtonian dynamics. (iv) The deep MOND limit corresponds to $a_o \rightarrow \infty$; phenomenology points to $\vec{A} \propto a_o^{-1}$, in this limit. We do not know that this should hold universally (i.e. for all sort of trajectories etc.), we shall assume this behavior for circular orbits, and for general trajectories assume only $\vec{A} \rightarrow 0$. We also do not have observational information on the very limit $a_o \rightarrow \infty$; reliable information exists only down to $a/a_o \sim 0.1$. Here we shall assume this everywhere for large a_o , but most of what we say is independent of this assumption. (v) We assume a_o to be time independent, and so we are free to assume that $\vec{A}$ does not depend explicitly on time. If a_o actually varies with cosmic time, our results may be considered the lowest order in an adiabatic approximation. Explicit time dependence could also enter $\vec{A}$ if the particle has variable mass, which is simple to include (but which we bar). (vi) $\vec{A}$ is invariant under translations, and transforms as a vector under rotations of the trajectory. (vii) We do insist on full Galilei invariance.

Indication that we need require some invariance connected with a velocity boost could have come from the study of unbound trajectories. When a fast, unbound particle of velocity v scatters gravitationally off some mass, with pericenter distance r , the characteristic acceleration around pericenter is $a \approx v_{es}^2/r$, where v_{es} is the escape speed from r . We can, however, with the aid of v , construct other quantities with the dimensions of acceleration, e.g. $a_n = (v^n \frac{d^n}{dt^n} a)^{1/(n+1)}$. Around pericenter, $a_n \approx (v/v_{es})^{2n/(n+1)} a$, which may be much larger than a itself. This might lead, in theories such as we discuss here, to much less of a mass discrepancy for such unbound particles, compared with that manifested by bound particles; for the latter, all quantities of the dimensions of acceleration are of order a . It is the appearance of the velocity v that engenders such terms. Galilei invariance would bar the appearance of v in the equation of motion, and all the permissible quantities with dimensions of acceleration are of the order of the acceleration, near pericenter. We may then

expect that bound, and scattering test particles will point to mass discrepancies of the same order (a_o/a). Observations of lensing by galaxy clusters may illuminate this question (for these v/v_{es} is typically a few hundred), pending the understanding of the relativistic extension of MOND.

While, for convenience, we treat motion in a static potential, all our results apply to the relative motion in the center-of-mass frame, of a system of two particles, interacting through a potential that depends on the relative position. So Galilei invariance of $\vec{A}$ for the particles assures invariance for the system.

Equation (3) satisfies all the above requirements, but does not conform to another the we now add; viz., that the equation of motion be derivable from an action principle. As far as we can see, there is nothing amiss, in principle, with effective equations of motion that are not so derivable. If their symmetries do not lead to conservation laws, this may just reflect interaction with the hidden degrees of freedom. In fact, exactly such effective theories have been proposed as models for dissipative systems[28]. That being as it may, our purpose here is to investigate equations of motion with an action underpinning. This is a strong constraint that enables us to say much about the theory. One result of this assumption is that a theory whose kinetic action is Galilei invariant, and has the correct Newtonian limit, and the required MOND limit, cannot be local; i.e. $\vec{A}$ in eq. (10) cannot be a function of a finite number of time derivatives of $\vec{r}(t)$. (From this we understand why eq. (3), which is local, cannot be derived from an action.) Other results concern a virial relation, conservation laws, and adiabatic invariants. In yet another we shall show that circular orbits are necessarily solutions of the equation of motion in a potential with the appropriate symmetry, and derive explicit expressions for the rotational speed.

We formulate the action principle as follows: There exists a kinetic action functional $S_k[\vec{r}(t), t_1, t_2, a_o]$ (not necessarily Lagrangian) that is a functional of the trajectory, $\vec{r}(t)$, given between two end times t_1 , and t_2 . Further assume that the variation of S_k under a change $\delta\vec{r}(t)$ may be written as

$$\delta S_k = -T^{-1} \int_{t_1}^{t_2} \vec{A}(t) \cdot \delta\vec{r}(t) dt + T^{-1} [\vec{\mathfrak{S}} \cdot \delta\vec{r}(t)]_{t_1}^{t_2}, \quad (11)$$

where $\vec{A}$ does not depend on the end points. The second term on the right-hand side is (up to the T normalization) the difference between the values of $\vec{\mathfrak{S}} \cdot \delta\vec{r}(t)$, at the two end times; $\vec{\mathfrak{S}}$ is some linear vector operator, acting on functions of t , [and may depend on $\vec{r}(t)$]. We then consider only equations of motion (10) whose kinetic part $\vec{A}$ can be obtained in this way from an action functional. The normalization by the time interval $T \equiv t_2 - t_1$ is introduced for convenience, so as to render the action finite when $t \rightarrow \infty$; it does not affect the equation of motion.

All the above requirements we made of $\vec{A}$ are now carried, *mutatis mutandis*, to the kinetic action S_k .

Because we want the kinetic part of the equation of motion to bear the modification that MOND entails, we choose to discuss the kinetic part separately. The full action of the theory is

$$S = S_k + S_p, \quad (12)$$

where the potential part has the usual form (except for our normalization):

$$S_p = -T^{-1} \int_{t_1}^{t_2} \phi \, dt. \quad (13)$$

The variation of the full action is

$$\delta S = -T^{-1} \int_{t_1}^{t_2} [\vec{A}(t) + \vec{\nabla}\phi] \cdot \delta\vec{r}(t) \, dt + T^{-1} [\vec{\mathfrak{S}} \cdot \delta\vec{r}(t)]_{t_1}^{t_2}. \quad (14)$$

Note that we do not define the physical trajectories as extrema of the action; that would be the case only if one restricts himself to variation, $\delta\vec{r}(t)$, for which the ends term vanishes, which is usually done by fiat. The trajectory may well be an extremum, but we need not and wish not make such requirements.

Standard Lagrangian theories are special instances (see below), but the above definition will suit us in more general cases. When the theory is non-local, it usually makes sense only when we take t_1 to $-\infty$, and t_2 to $+\infty$. In this case the equation of motion is a non-local equation for the full world-line of the particle. A finite trajectory (specified between finite times) is permitted if it is a segment of a world-line that solves the equation of motion. An important question regarding a given theory is how many (vector) parameters are needed to specify a permitted world line; this is also the number of initial conditions that one has to specify for a finite time t .

Consider now variations of S under coordinate transformations. For solutions of the equation of motion, only the ends term remains in eq. (14), and it is thus clear that $\vec{\mathfrak{S}}$ generates conserved quantities, when underlying symmetries of S are present. For example, if S is invariant under translations $\vec{r} \rightarrow \vec{r} + \delta\vec{\epsilon}$, we see that the difference of the values of $\vec{\mathfrak{S}}(1)$ ($\vec{\mathfrak{S}}$ acting on 1) between the two end points must vanish. This quantity may be identified as the momentum. Similarly, if S is invariant under rotations about a direction $\vec{n}$, since $\delta\vec{r} = \epsilon\vec{n} \times \vec{r}$, we find that $-\vec{n} \cdot [\vec{\mathfrak{S}} \times \vec{r}(t)]$ is the same at the two end points. Thus we identify $-\vec{\mathfrak{S}} \times \vec{r}(t)$ with the angular momentum.

Now take a dilatational increment in a solution of the equation of motion: $\delta\vec{r}(t) = \epsilon\vec{r}(t)$. For trajectories on which $\vec{\mathfrak{S}} \cdot \vec{r}(t)$ does not increase as fast as t , at $t \rightarrow \pm\infty$ we get, from eq. (11), for the variation of S_k (for the full world line $T \rightarrow \infty$)

$$\delta S_k = -\epsilon \langle \vec{A} \cdot \vec{r}_p \rangle, \quad (15)$$

where $\langle \rangle$ is the time average over the trajectory. This certainly holds for periodic trajectories, but surmisedly for any bounded trajectory. We now calculate δS_k differently: On dimensional grounds

$$S_k[\lambda \vec{r}(t), \lambda a_o] = \lambda^2 S_k[\vec{r}(t), a_o], \quad (16)$$

for a constant λ . Putting $\lambda = 1 + \epsilon$, and identifying the first order terms in ϵ , we get, as in Euler's theorem for homogeneous functions,

$$\delta S_k + \epsilon a_o \frac{\partial S_k}{\partial a_o} = 2\epsilon S_k. \quad (17)$$

Comparing with expression (15) for δS_k we get

$$-\langle \vec{A} \cdot \vec{r} \rangle = 2S_k - a_o \frac{\partial S_k}{\partial a_o}, \quad (18)$$

and from the equation of motion

$$S_k(1 + \frac{1}{2} \hat{S}_k) = \frac{1}{2} \langle \vec{r} \cdot \vec{\nabla} \phi \rangle, \quad (19)$$

where $\hat{S}_k \equiv -\frac{\partial \ln S_k}{\partial \ln a_o}$. This is a generalization of the virial relation (in the standard Newtonian case S_k is the mean kinetic energy, with our normalization).

More generally, if S_k depends on a number of dimensional constants, C_ℓ , $\ell = 1, \dots, I$, with dimensions $(length)^{\alpha_\ell} \times (time)^{\beta_\ell}$, respectively, then $\hat{S}_k$ in eq. (19) is replaced by $-\sum_{\ell=1}^I \alpha_\ell \frac{\partial \ln S_k}{\partial \ln C_\ell}$. (If we replace S_k with S , and put $\phi = 0$ we get a more general relation that reduces to the former when $S = S_k + \phi$ with ϕ depending only on $\vec{r}$, because then we must have $\phi = C^2 f(\vec{r}/r_0)$ with C and r_0 of dimensions of velocity and length, respectively.)

In Newtonian dynamics $\vec{A} = \vec{a}$ is linear in $\vec{r}(t)$. In a MOND theory this may not be: From dimensional arguments

$$\vec{A}[\lambda \vec{r}(t), \lambda a_o] = \lambda \vec{A}[\vec{r}(t), a_o], \quad (20)$$

or, taking $\lambda = a_o^{-1}$,

$$\vec{A}[\vec{r}(t), a_o] = a_o \vec{A}[a_o^{-1} \vec{r}(t), 1]. \quad (21)$$

If $\vec{A}$ is linear in $\vec{r}(t)$, it is independent of a_o altogether.

It may be more convenient, sometimes, to write the equation of motion in terms of the expansion coefficients of $\vec{r}(t)$ in some basis of complete functions, instead of $\vec{r}(t)$ (which is itself the expansion coefficient of the trajectory in the functions $\delta(t' - t)$). This is not a common practice because (i) for time-local, Lagrangian theories the locality is manifest only when t is used as an independent variable (ii) t is a natural choice when the potential is time independent.

If we forgo locality anyway, the benefits of using a different decomposition of the trajectory may outweigh the drawbacks. For example, we may use the Fourier transform, $\vec{\rho}(\omega)$, of $\vec{r}(t)$. As a functional of $\vec{\rho}$ the kinetic action may take a particularly simple form. We discuss examples below.

It is well known that theories with higher derivatives tend to suffer from various diseases. we discuss these and their prevention at the end of this section, after we have considered a few examples.

LAGRANGIAN THEORIES

We now specialize to kinetic actions that are derivable from a Lagrangian function:

$$S_k = T^{-1} \int_{t_1}^{t_2} L_k dt \quad (22)$$

(the total Lagrangian being $L = L_k - \phi$). We assume ϕ to depend only on $\vec{r}$ (and, in particular, not to depend explicitly on the time). In this case $\vec{A}$ is given by the standard Euler-Lagrange expression

$$\vec{A} = \sum_{i=1}^N (-1)^{i-1} \left[\frac{\partial L_k}{\partial \vec{r}^{(i)}} \right]^{(i)}, \quad (23)$$

and the operator $\vec{\mathfrak{S}}$ is

$$\vec{\mathfrak{S}} = \sum_{m=1}^N \vec{p}_m D^{m-1}, \quad (24)$$

with

$$\vec{p}_m \equiv \sum_{i=m}^N (-1)^{i-m} \left[\frac{\partial L_k}{\partial \vec{r}^{(i)}} \right]^{(i-m)}. \quad (25)$$

Here, $D \equiv \frac{d}{dt}$, and a superscript (i) stands for the i th time derivative.

If the number, N , of derivatives on which L_k depends is finite, the theory is local. Even when it is non-local, the theory may be a limit of a sequence of local ones with Lagrangians L_k^N , $N \rightarrow \infty$. (A common example is an L_k that can be expanded in a power series in some parameter— a_o in our case—where the coefficients are function of a finite number, N , of derivatives.) When it is not even that we call the theory strongly non-local. For such theories the sums in the expressions above do not converge for at least some trajectories. These expressions may still be of value, as they may converge for some trajectories, and may be analytically continued for even more trajectories. A different expression for $\vec{A}$ must then be given, to establish an equation of motion.

As explained above, translation invariance implies the conservation of the momentum $\vec{\mathfrak{S}}(1)$, which here is just $\vec{p}_1$ (this can also be seen from the fact that $\vec{A} = \frac{d\vec{p}_1}{dt}$). (In a many-body system with bodies of masses m_i , and potentials that depend on coordinate differences, $\sum_i m_i \vec{A}_i = 0$.) Similarly, the angular

momentum—whose component along $\vec{n}$ is conserved when ϕ is symmetric under rotations about $\vec{n}$ —is given by

$$\vec{J} = -\vec{\mathfrak{S}} \times \vec{r}(t) = \sum_{m=1}^N \vec{r}^{(m-1)} \times \vec{p}_m = \sum_{m=1}^N \vec{r}^{(m-1)} \times \sum_{i=m}^N (-1)^{i-m} \left[\frac{\partial L_k}{\partial \vec{r}^{(i)}} \right]^{(i-m)}, \quad (26)$$

or

$$\vec{J} = \sum_{i=1}^N \sum_{m=1}^i (-1)^{i-m} \vec{r}^{(m-1)} \times \left[\frac{\partial L_k}{\partial \vec{r}^{(i)}} \right]^{(i-m)}. \quad (27)$$

This also follows straightforwardly from the fact that $\vec{r} \times \vec{A} = \frac{d\vec{J}}{dt}$, so the rotational symmetry of ϕ [which reads $\vec{\nabla} \phi \cdot (\vec{n} \times \vec{r}) = 0$] implies, from the equation of motion, $\frac{d(\vec{n} \cdot \vec{J})}{dt} = 0$.

As in standard dynamics, if the potential is rotationally symmetric about $\vec{n}$, $\vec{n} \cdot \vec{J}$ is conserved, even when the Lagrangian depends explicitly on time, such as when a_o varies with cosmological time. Obviously, this has implications for galaxy evolution that is driven by such variations.

If L does not depend explicitly on time, the energy H is conserved, where

$$H = \sum_{i=1}^N \sum_{m=1}^i (-1)^{m-1} \vec{r}^{(i-m+1)} \cdot \left[\frac{\partial L}{\partial \vec{r}^{(i)}} \right]^{(m-1)} - L. \quad (28)$$

We separate $H = H_k + \phi$, and call H_k the kinetic energy. The conservation of H can be deduced by noting that $\vec{v} \cdot \vec{A} = \dot{H}_k + \frac{\partial L_k}{\partial t}$, so that, from the equation of motion, if $\frac{\partial L_k}{\partial t} = 0$, we have $\dot{H}_k + \dot{\phi} = 0$.

If we add to a total derivative to L , then $\vec{A}$, H , p_1 , and $\vec{J}$ remain the same, but p_i for $i > 1$, and hence $\vec{\mathfrak{S}}$ do change.

In appendix B we derive an expression for the energy in general Lagrangian theories, which, specialized to the case of MOND, reads

$$H = L_k - 2a_o \frac{\partial L_k}{\partial a_o} + \phi + \dot{Q}, \quad (29)$$

where

$$Q = - \sum_{i=2}^N \sum_{m=1}^{i-1} (-1)^{m-1} (i-m) \vec{r}^{(i-m)} \cdot \left[\frac{\partial L_k}{\partial \vec{r}^{(i)}} \right]^{(m-1)}. \quad (30)$$

This has an immediate corollary: for any solution of the equation of motion on which Q does not grow as fast as t , at $t \rightarrow \pm\infty$, such as on periodic, and presumably other bounded trajectories, we can write

$$H = \langle H \rangle = \langle L_k \rangle - 2a_o \frac{\partial \langle L_k \rangle}{\partial a_o} + \langle \phi \rangle. \quad (31)$$

Combined with the virial relation (19) this gives

$$\langle H_k \rangle = \frac{1}{2} \gamma \langle \vec{r} \cdot \vec{\nabla} \phi \rangle, \quad (32)$$

where $\gamma \equiv (1 + 2\hat{S}_k)/(1 + \hat{S}_k/2)$. In the Newtonian limit $\gamma = 1$; in general $\gamma < 4$; for circular orbits in the MOND limit $\gamma = 2$.

If the theory is Galilei invariant and is local or weakly non-local, L_k may be written (up to a total time derivative) in the form

$$L_k = \frac{1}{2} \alpha v^2 + \tilde{L}_k(a_o, \vec{r}^{(2)}, \vec{r}^{(3)}, \dots), \quad (33)$$

where α is a number, and $\tilde{L}_k$ depends only on the acceleration and higher derivatives of $\vec{r}(t)$. We show this in appendix A. Correspondence with Newtonian dynamics requires that $\alpha = 1$, and that $\tilde{L}_k$ vanish for $a_o = 0$ (or reduce to a time derivative, which we can eliminate from the outset). On the other hand, the MOND limit requires $\alpha = 0$. Thus, *Candidate, Galilei-invariant theories for MOND, that are derivable from an action, must be strongly non-local.* This is the basis for our statement that eq. (3), which is local, and Galilei invariant, cannot be derived from an action.

Higher-derivative Lagrangian theories that are not strongly non-local may be cast in a Hamiltonian form as described by Ostrogradski[29]. Very succinctly, one defines N coordinates, the first of which is $\vec{q}_1 \equiv \vec{r}$; the rest are defined recursively by $\vec{q}_m = \frac{d\vec{q}_{m-1}}{dt}$, $m = 2, \dots, N$. There are N momenta, which are those defined by eq. (25). The Hamiltonian is H , as defined in (28), in which the derivatives of $\vec{r}(t)$ have been expressed by the $2N$ coordinates and momenta, taken to be independent. The Hamilton equations of the usual form then give the equation of motion, and, to boot, the above relations between the coordinates. The Hamiltonian formalism is particularly useful when one endeavors to quantize the theory; we shall have little use for it here.

SOME HEURISTIC MODEL ACTIONS

Before embarking on general issues concerning properties of the solutions, we describe a few examples of theories on which we shall demonstrate various points.

Consider first a class of kinetic Lagrangians of the form

$$L_k = \frac{1}{2} \vec{v} f(a_o^2 \overleftarrow{D} g^{-1} \overrightarrow{D}) \vec{v}. \quad (34)$$

Here, $\overleftarrow{D}$ and $\overrightarrow{D}$ are the time derivative operators acting to their left and right respectively, and g is a function of the derivatives of $\vec{r}(t)$ from the second upward, having the right dimensions (l^2/t^6) to make the argument of f dimensionless. We must have $f(0) = 1$ for Newtonian correspondence; furthermore, as stated in point (iii) in the beginning of this section, for L_k to approach the

Newtonian limit for all trajectories, $f(z)$ has to be non-singular at $z = 0$ in the complex plane. If it has a Taylor expansion, with a non-zero radius of convergence:

$$f(z) = 1 + \sum_{m=1}^{\infty} b_m z^m, \quad (35)$$

then

$$L_k = \frac{1}{2}v^2 + \frac{1}{2}b_1 a_o^2 g^{-1} a^2 + \frac{1}{2}b_2 a_o^4 \left[\frac{d(g^{-1}\vec{a})}{dt} \right]^2 + O(a_o^6), \quad (36)$$

which has the form (33). Clearly, L_k may not be represented by this expansion for all values of a_o , lest it not be strongly non-local, and not have a MOND limit. In the next section we derive an exact expression for the rotation curve in this class of theories, and show that to obtain asymptotically flat rotation curves we must have $f(x) \propto x^{-1/2}$ for $x \rightarrow \infty$ (x real). For example $f(z) = (1+z)^{-1/2}$ satisfies all the above requirements; it has a finite radius of convergence.

The Lagrangian (34) is Galilei invariant: the required Newtonian limit implies that $f(a_o^2 \overleftarrow{\overrightarrow{D}} g^{-1} \overrightarrow{\overleftarrow{D}})$ acting on a constant vector $\vec{v}_0$, to its left or right, gives $\vec{v}_0$ itself; so, under a transformation $\vec{r}(t) \rightarrow \vec{r}(t) + \vec{v}_0 t + \vec{r}_0$, L_k changes by $v_0^2/2 + \vec{v}_0 \cdot \vec{v}(t)$, which is a total time derivative.

Second, consider Lagrangians of the form

$$L_k = \frac{1}{2} \vec{v} f[a_o^2 (\overleftarrow{\overrightarrow{D}} h \overrightarrow{\overleftarrow{D}})^{-1}] \vec{v}. \quad (37)$$

Here, Galilei invariance is achieved because of the required MOND limit $f(\infty) = 0$.

We consider also actions of the form (22), where L_k depends also on parameters, g_i , that are functionals of the whole trajectory (and not functions of the local values of $\vec{r}(t)$, and its derivatives). Take, for instance L_k of the form (34) [or (37)] with g (or h) of the form

$$g = \langle G \rangle \equiv T^{-1} \int G dt, \quad (38)$$

where G is a function of the derivatives of $\vec{r}(t)$. The action is not, strictly speaking, Lagrangian; we keep that epithet but call such Lagrangians compound. When all the g_i s are of the form $g_i = \langle G_i \rangle$, with local G_i s, the equation of motion has the same form as the Euler-Lagrange equation, with L_k replaced by

$$M_k = L_k + \sum_i G_i \frac{\partial L_k}{\partial g_i}. \quad (39)$$

In an even wildier class of (non-Lagrangian) theories we define the kinetic action S_k as a solution of the equation

$$S_k = \frac{1}{2} T^{-1} \int \vec{v} \cdot F \left(\frac{S_k \overleftarrow{\overrightarrow{D}} \overrightarrow{\overleftarrow{D}}}{a_o^2} \right) \vec{v} dt. \quad (40)$$

The integrand is defined when $\vec{v}(t)$ can be written as a sum of exponentials $\exp(\alpha t)$, with arbitrary complex α : $e^{\alpha t} F\left(\frac{S_k \overleftrightarrow{DD}}{a_o^2}\right) e^{\beta t} = F\left(\frac{S_k \alpha \beta}{a_o^2}\right) e^{(\alpha+\beta)t}$. In the Newtonian limit $F(x) \xrightarrow{x \rightarrow \infty} 1$. To get the correct MOND limit for circular orbits, we must have $F(x) = (x/\eta)^{1/3}$ for $x \ll 1$ (with our normalization of a_o , $\eta = 9/8$). $F(z)$ is analytic at infinity, according to our general requirement, but is non-analytic at $z = 0$. This theory is “almost” linear, whence come its heuristic value.

Note that only trajectories that can be expanded in real frequencies are relevant here: For an infinite-time world line there is no finite- S_k solution to eq. (40), if there are complex-frequency components to $\vec{r}(t)$. Such orbits necessarily approach the Newtonian regime, for large (positive or negative) times, but they are irrelevant there as candidate solutions of the equation of motion.

The equation of motion is

$$(1 + \hat{S}_k/2) F\left(-\frac{S_k D^2}{a_o^2}\right) D^2 \vec{r}(t) = -\vec{\nabla} \phi \quad (41)$$

with $\hat{S}_k \equiv -\frac{\partial \ln S_k}{\partial \ln a_o}$.

To obtain eq. (41) we write

$$\begin{aligned} \delta S_k = \frac{1}{2T} \int dt \left[\frac{d\delta \vec{r}(t)}{dt} \cdot F\left(\frac{S_k \overleftrightarrow{DD}}{a_o^2}\right) \vec{v}(t) + \vec{v}(t) \cdot F\left(\frac{S_k \overleftrightarrow{DD}}{a_o^2}\right) \frac{d\delta \vec{r}(t)}{dt} \right] \\ + \delta S_k \frac{1}{2T} \int dt \vec{v}(t) F'\left(\frac{S_k \overleftrightarrow{DD}}{a_o^2}\right) \frac{\overleftrightarrow{DD}}{a_o^2} \vec{v}(t). \end{aligned} \quad (42)$$

The two terms in the first integral are equal, as $\overleftrightarrow{DD}$ acts symmetrically. The second integral can be written in terms of $\hat{S}_k$, leading to

$$\delta S_k = (1 + \hat{S}_k/2) \frac{1}{T} \int dt \frac{d\delta \vec{r}(t)}{dt} \cdot F\left(\frac{S_k \overleftrightarrow{DD}}{a_o^2}\right) \vec{v}(t). \quad (43)$$

Now, when $\vec{r}(t)$ and $\delta \vec{r}(t)$ can be expanded in Fourier components with real frequencies, the last integral equals $-\int dt \delta \vec{r}(t) \cdot F\left(-\frac{S_k D^2}{a_o^2}\right) D^2 \vec{r}(t)$, up to end terms [when $F(z)$ is analytic at $z = 0$ this is seen by integration by parts; when we go via Fourier space the end terms are lost], from which follows eq. (41).

To insure Galilei invariance, L_k must actually be invariant, not just incremented by a time derivative; S_k itself must be invariant—not just changed by end terms—as can be seen from the equation of motion. This, in turn, is insured by the required MOND limit [$F(0) = 0$]. (Hence the choice of $\overleftrightarrow{DD}$

in the argument of F , in the definition of S_k , is essential. A choice of $-D^2$ instead would have given an expression that differs by an ends term, and that is not exactly invariant under Galilei transformations.) We want S_k to be determined uniquely by eq. (40). If this has many solutions, for a given trajectory, we need an added prescription to choose from among them. The obvious desideratum that S_k go to its Newtonian value, for that trajectory, when $a_o \rightarrow 0$ may do (see next subsection); it eliminates $S_k = 0$, which is always a solution. Note that the theory is time-reversal symmetric (if S_k is a solution so is its value for the time reversed orbit, and the two have the same Newtonian limit).

Finally, we bring for comparison a local Lagrangian theory with

$$L_k = \frac{1}{2}v^2 f(a_o^2/a^2); \quad (44)$$

obviously it not Galilei invariant.

THE NATURE OF THE SOLUTIONS OF NONLOCAL THEORIES

Higher-derivative theories are notoriously problematic. Pais and Uhlenbeck[30], and more recently, Jaén et al.[31], Eliezer and Woodard[32], and Simon[33] review these problems in Lagrangian theories, describe possible solutions, and give further references. Very succinctly, the problems are: (i) The energy as defined in eq. (28) is not bound from below: there are solutions of the equation of motion with arbitrarily small (large negative) energy. (ii) There appear runaway solutions of the equation of motion, on which the velocity, acceleration, etc. blow up at $t \rightarrow \infty$. In linear theories the divergence is exponential. (iii) When N is the highest derivative on which the Lagrangian depends, the equation of motion is of order $2N$, and require the specification of that many initial conditions. While this may not be a matter-of-principle problem, it is disconcerting. There is a connection between these three problems, and they can usually be eliminated in one fell swoop, by adding appropriate constraints to a finite order theory, or by culling out the majority of the solutions of the equation of motion that are not analytic in some expansion parameter[31,32,33].

We can see the relation between the runaway character of a solution and the energy from eq. (31): When the solution is not runaway, the assumption underlying this relation is valid, and then we see that H is bound from below, since L_k , and $-\frac{\partial L_k}{\partial a_o}$ are positive definite.

When the theory is of infinite order, we may expect exacerbation of the above problems. For instance, one may have to dictate an infinite number of initial conditions, and this may signify a total loss of the initial-value formulation[32].

Can MOND-oriented theories avoid these daunting problems? As has been emphasized by Eliezer and Woodard[32] a strongly non-local theory (in our terminology) is not a higher-derivative theory, and need not suffer from any of the problems endemic to such theories. We saw that Galilei-invariant

MOND theories must be strongly non-local and may well be trouble free. Theories that emerge as effective approximation to healthy theories, by an elimination of some degrees of freedom, are obviously as healthy as their progenitors. As explained in Sec. II we envisage MOND to come about in just this way.

We now demonstrate that the theory whose action is defined by eq. (40), with a (solvable) harmonic-oscillator potential $\phi = \frac{1}{2}\omega_0^2 r^2$, is free of the above problems. As explained below eq. (40), only trajectories that can be expanded in Fourier components with real frequencies have a chance of solving the equation of motion (41). From this equation we also see that every frequency ω that appears in the expansion must satisfy

$$(1 + \hat{S}_k/2)F\left(\frac{S_k\omega^2}{a_o^2}\right)\omega^2 = \omega_0^2. \quad (45)$$

We can choose F to give a unique (positive) solution for ω^2 , given S_k . The physical trajectories have thus a single frequency, and can be written as

$$\vec{r}(t) = \frac{1}{\sqrt{2}}(\vec{r}_0 e^{i\omega t} + \vec{r}_0^* e^{-i\omega t}). \quad (46)$$

The defining equation of S_k , (40), which supplements the equation of motion, reads here

$$S_k = \frac{1}{2}\omega^2 r_0^2 F\left(\frac{S_k\omega^2}{a_o^2}\right), \quad (47)$$

where $r_0^2 \equiv \vec{r}_0 \cdot \vec{r}_0^*$. We can also write from eqs. (45) (47)

$$(1 + \hat{S}_k/2)S_k = \frac{1}{2}r_0^2\omega_0^2. \quad (48)$$

Take for instance

$$F(x) = (1 + \eta x^{-1})^{-1/3}, \quad (49)$$

which has the desired Newtonian and MOND limits (with $\eta = 9/8$). It is easy to ascertain that $Re[\vec{r}_0]$ and $Im[\vec{r}_0]$ determine ω^2 (and S_k) uniquely.

Thus, there are no runaway solutions, and the physical trajectories are determined by two initial conditions, just as for a Newtonian harmonic oscillator. Here too the motion is harmonic, but the frequency depends on the amplitude. For example, in the deep MOND case—i.e., when $\omega_0^2 r_0 \ll a_o$, we have $\hat{S}_k \approx 1$, and $F(x) \approx (x/\eta)^{1/3}$, and ω is determined by

$$\omega^2 \approx \omega_0^2 \left(\frac{\omega_0^4 r_0^2}{a_o^2}\right)^{-1/4} \gg \omega_0^2. \quad (50)$$

A circular orbit is a special case which we shall discuss below.

The case of an anisotropic harmonic potential, which is as easy to solve, is also instructive. The motion is still harmonic with a unique frequency in

each of the principal directions of the potential, the equation of motion now being

$$(1 + \hat{S}_k/2)F\left(\frac{S_k\omega_m^2}{a_o^2}\right)\omega_m^2 = \omega_{0m}^2, \quad m = 1, 2, 3. \quad (51)$$

with

$$(1 + \hat{S}_k/2)S_k = \sum_{m=1}^3 |r_{0m}|^2 \omega_{0m}^2. \quad (52)$$

Given the three amplitudes and phases encoded in the complex amplitudes r_{0m} one determines S_k and the three frequencies ω_m ; each depends not only on its own amplitude; they are coupled through S_k .

In general, exponentially exploding solutions are precluded in MOND, because they possess derivatives that all blow up at infinite time, and this necessarily takes us to the Newtonian regime where they are not physical trajectories. In finite-order MOND theories even trajectories that runaway as a high enough power of t are precluded. This, in itself, does not solve any of the problems listed above. We saw that the criterion for a non-runaway solution is that Q of eq. (30) not grow as fast as t at $\pm\infty$. Take, for instance, the local MOND Lagrangian given by eq. (44). Clearly, the equation of motion cannot have solutions for which a increases indefinitely at $t \rightarrow \infty$; still, the theory has all the usual illnesses: The runaway solutions are such that $v \propto t^{1/3}$ for $t \rightarrow \infty$, so $a \rightarrow 0$, but the $\dot{Q}$ term in the expression for H is $\propto v^3$, enough to qualify as a runaway.

IV. CIRCULAR ORBITS: DYNAMICS OF AXISYMMETRIC DISK GALAXIES

No other observation is as central to the discussion of the mass discrepancy in galactic systems, and as clear-cut, as the measurement and analysis of rotation curves of disk galaxies. Fortunately, the relevant trajectories are particularly amenable to treatment in the framework that we consider here. These are circular, constant-speed orbits (the latter epithet will be implicit hereafter) in the mid-plane of an axisymmetric, and plane symmetric potential field, ϕ .

THE EQUATION OF MOTION-ROTATION CURVES

We now show that, under the MOND assumption, and for a general action of the type given by eqs. (11) - (13), the speed, v , on a circular orbit, in the mid-plane of a potential with the symmetry of a disk, depends on the orbital radius, r , through a relation of the form

$$\mu(a/a_o)a = \frac{\partial\phi}{\partial r}, \quad (53)$$

where $\mu(x)$ depends on the specific kinetic action, and $a = v^2/r$. Thus, eq. (3) gives the exact expression for the rotation curves, and existing rotation-curve analyses, which have all employed this expression[2,4,5,7], are valid in the present framework.

For consistency we first ascertain that, for a circular orbit, the functional derivative, $\vec{A}$, in the equation of motion, is a radial vector of a constant magnitude. This is a property of the kinetic action alone (irrespective of the potential) and is necessary if circular orbits are to be solutions of the equation of motion in the mid-plane. This property follows from the assumed translation and rotation invariances of the kinetic action. Under a infinitesimal translation $\vec{r} \rightarrow \vec{r} + \vec{\epsilon}$, S_k changes by $\vec{\epsilon} \cdot \int \vec{A} dt$, according to eq. (11) for the variation of S_k [the ends term vanishes as $\vec{r}(t)$ and $\delta\vec{r}(t)$ are periodic]. This must vanish for arbitrary $\vec{\epsilon}$, so $\int \vec{A} dt = 0$. From the symmetry of the orbit, the component of $\vec{A}$ perpendicular to the orbit must be constant and hence must vanish. Similarly, the invariance of S_k under a rotation about an arbitrary axis $\vec{n}$ ($\delta\vec{r} \propto \vec{n} \times \vec{r}$) implies that $\int \vec{A} \times \vec{r}(t) dt = 0$. Thus, $\vec{A}$ cannot have a tangential component either, and is a radial vector of constant magnitude, as required. The action underpinnings is essential; it is easy to write an equation of motion (10) in which $\vec{A}$ has the above symmetries but is not radial for circular orbits. (But TR is then not a symmetry.)

The magnitude of $\vec{A}$ depends only on the parameters of the orbit, to wit v , and r . Because A/a is dimensionless, it must, by the MOND assumption, be of the form $\mu(a/a_o)$, leading to

$$\vec{A} = \mu(a/a_o)\vec{a}, \quad (54)$$

and to eq. (53), in turn. For there to be a circular-orbit solution for every positive value of $\frac{\partial\phi}{\partial r}$, $x\mu(x)$ must take all positive values, and $\vec{A}$ must be directed inward. Uniqueness requires that $x\mu(x)$ is monotonic. As a corollary we get that any kinetic action that is rotation and translation invariant necessarily gives the Newtonian relation for circular orbits, if it does not contain any dimensional constant.

There exists a simple and useful expression for $\mu(x)$ in terms of the values of the action for circular orbits:

$$\mu(a/a_o) = 2v^{-2}S_k^c(1 + \frac{1}{2}\hat{S}_k^c), \quad (55)$$

where S_k^c is the action calculated for a circular orbit, and $\hat{S}_k^c = -\frac{\partial \ln S_k^c}{\partial \ln a_o}$. This relation enables us to circumvent the calculation of functional derivatives.

Equation (55) follows from the general expression (18) for $\langle \vec{A} \cdot \vec{r} \rangle$ —which holds for any periodic orbit—with eq. (54), and the equality $\vec{a} \cdot \vec{r} = -v^2$.

With our normalization, S_k has the dimensions of velocity square; so, S_k^c must be of the form

$$S_k^c = \frac{1}{2}v^2\lambda(a/a_o). \quad (56)$$

(Any functional of the orbit reduces to a function of v , r , and a_o , for a circular orbit. Thus, S_k^c/v^2 is a dimensionless function of these, and must be of the form $\frac{1}{2}\lambda(v^2/ra_o)$.) Hence,

$$\mu(x) = \lambda(x) + \frac{1}{2}x \frac{d\lambda}{dx} = \lambda(x)[1 + \hat{\lambda}(x)/2], \quad (57)$$

where $\hat{\lambda}(x) \equiv d \ln \lambda / d \ln x$.

Only the time-reversal-symmetric part of S_k contributes to S_k^c , and hence to μ , because, for a circular orbit, time reversal is equivalent to some π -rotation about an axis in the orbital plane, under which S_k is invariant.

When the action is derivable from a Lagrangian, L_k -simple or compound, local or not—that does not depend on the time explicitly, one has $S_k^c = L_k^c$, where L_k^c is the constant value of the Lagrangian on the circular orbit. Then

$$\mu(a/a_o) = 2v^{-2}L_k^c(1 + \frac{1}{2}\hat{L}_k^c), \quad (58)$$

with $\hat{L}_k^c = -\frac{\partial \ln L_k^c}{\partial \ln a_o}$.

Consider, for instance, Lagrangian actions with L_k of the form (34). The function g is constant on a circular orbit, and, in fact, must be proportional to $r^{-4}v^6$, on dimensional grounds. The proportionality constant may be absorbed into a_o . The operator $\overleftrightarrow{D}D$, when sandwiched between any two derivatives of $\vec{r}_c(t)$ (or $-D^2$, when acting on any of these), is just the multiplication by $\omega^2 = v^2/r^2$. We then find that

$$L_k^c = \frac{1}{2}v^2 f(a_o^2/a^2), \quad (59)$$

from which results the expression

$$\mu(a/a_o) = f(x)[1 + \hat{f}(x)]|_{x=a_o^2/a^2}, \quad (60)$$

where $\hat{f}(x) \equiv -d \ln f / d \ln x$.

Since μ does not depend on the choice of g , many theories give the same rotation curve for a given mass distribution. For similar reasons, Lagrangians of the form (37) give the same rotation curve (with the same choice of f), although they are very different theories. The same is true of the compound Lagrangians discussed below eq. (37), and of the local Lagrangian given by eq. (44).

For actions of the form (40), we find from the general expression for $\mu(x)$ eq. (55) [or directly from the equation of motion (41)] that $\mu(x) = [1 + \frac{1}{2}\hat{\lambda}(x)]\lambda(x)$, where $\lambda(x)$ is the solution of $\lambda(x) = F[x^2\lambda(x)/2]$ with the correct Newtonian limit; i.e., the solution, for $\alpha = 0$, of $\lambda_\alpha(x) = F[x^2\alpha\lambda_\alpha(x)/2]$ that satisfies $\lambda_\alpha(x) \rightarrow 1$ for $\alpha \rightarrow \infty$.

Clearly, we can add to the Lagrangian a time-reversal-antisymmetric part without affecting the rotation curve. The same is true of a more general type of terms that vanish on circular orbits [e.g. $\propto (\vec{v} \cdot \vec{a})^2$].

The fact that the rotation curve, for a given mass distribution depends only on the value of the action for circular orbits is gratifying in that it allows us to test this general class of theories by rotation-curve analysis, without having to know the exact theory [by adopting a best value for a_o , and a

reasonable form for $\mu(x)$]. This, however, leaves us bereft of an important observational guide in discriminating between theories within the class.

THE ENERGY AND ANGULAR MOMENTUM

When the action is derivable from a Lagrangian, and energy and angular momentum are defined—in the sense that the formal sums that express them converge, or may be understood as analytic continuations—we find simple expressions for these quantities on circular orbits.

Because scalar functions are constant on circular orbits, $\dot{Q}$ in expression (29) for the energy vanishes and we have

$$H_c = L_k^c - 2a_o \frac{\partial L_k^c}{\partial a_o} + \phi. \quad (61)$$

In terms of $\lambda(x)$ of eq. (56), which now describes the dependence of L_k^c on a/a_o , we can write

$$H_c = \frac{1}{2} v^2 \nu(a/a_o) + \phi, \quad (62)$$

where

$$\nu(x) = \lambda(x)[1 + 2\hat{\lambda}(x)] = \mu(x)\kappa(x), \quad \kappa(x) = \frac{1 + 2\hat{\lambda}(x)}{1 + \hat{\lambda}(x)/2}. \quad (63)$$

In the deep MOND limit, where $\mu(x) \approx x$, and $\hat{\lambda}(x) \approx 1$, $\nu(x) \approx 2x$.

Putting together eqs. (62), (63), and eq. (53) [$\mu(a/a_o)a = \phi'$] which governs circular orbits in a disk potential ϕ (taken to vanish at infinity), we can write for such a motion

$$H_c = \frac{1}{2} \mu(a/a_o) v^2 (\kappa + 2/\hat{\phi}), \quad (64)$$

where $\hat{\phi} \equiv d \ln \phi / d \ln r$. In Newtonian dynamics ($\mu = 1$, $\kappa = 1$) there are no negative-energy circular orbits for potentials with $\hat{\phi} \leq -2$. It is interesting to note that in the strict MOND limit we have $\kappa \rightarrow 2$, and the limiting potential approaches the Newtonian or Coulomb potential, with $\hat{\phi} \rightarrow -1$.

Coming now to the angular momentum, we note that, for a circular orbit, vectors in the orbital plane [$\vec{r}(t)$ and its derivatives, derivative of L_k with respect to those, etc.] are radial or tangential, according to whether they are even or odd under time reversal, respectively—because, for such orbits, time reversal is equivalent to some π rotation about a diameter of the orbit. We see then from expression (27) for $\vec{J}$ that only the time-reversal-symmetric part of L_k contributes. Furthermore, the summand in eq. (27) can now be written (after successively shifting time derivatives from the second factor to the first) as $\vec{r}^{(m-1)} \times \frac{\partial L_k}{\partial \vec{r}^{(m)}}$. Hence, we can write for a circular orbit

$$\vec{J}_c = \sum_{m=1}^N m \vec{r}^{(m-1)} \times \frac{\partial L_k}{\partial \vec{r}^{(m)}}. \quad (65)$$

Since $\vec{r}^{(m-1)} = -\omega^{-2}\vec{\omega} \times \vec{r}^{(m)}$, with $\vec{\omega} = r^{-2}\vec{r} \times \vec{v}$, we have

$$\vec{J}_c = \omega^{-2}\vec{\omega} \sum_{m=1}^N m\vec{r}^{(m)} \cdot \frac{\partial L_k}{\partial \vec{r}^{(m)}}. \quad (66)$$

Finally, from expression (B-4) in appendix B, for this sum, we get

$$\vec{J}_c = \frac{2}{\omega^2}\vec{\omega} \left(L_k^c - a_o \frac{\partial L_k^c}{\partial a_o} \right). \quad (67)$$

In terms of $\lambda(x)$ of eq. (56) we have

$$\vec{J}_c = \eta(a/a_o)\vec{r} \times \vec{v}, \quad \eta(x) = \lambda(x)[1 + \hat{\lambda}(x)] = \mu(x) \frac{1 + \hat{\lambda}(x)}{1 + \hat{\lambda}(x)/2}. \quad (68)$$

So, in the very MOND limit $J_c \rightarrow 4v^3/3a_o$, and, interestingly, depends only on v .

In modifications of gravity, the expression for the angular momentum is the same as in conventional dynamics; so, the adiabatic invariance of $\vec{J}$ has rather different consequences in the two classes of theories based, respectively, on modifying gravity, and modifying the law of motion.

V. MODIFIED DYNAMICS AT LOW FREQUENCIES

Notwithstanding the phenomenological success of the MOND scheme, it is of interest to consider modifications that hinge on other system properties. After all, galactic systems differ from laboratory and planetary systems in attributes other than the typical accelerations in them. Here we examine modifications that hinge on the time scale involved. More accurately, we have in mind theories that involve a constant, Ω , with the dimensions of frequency. In analogy with the above discussion of MOND, the equation of motion is of the form

$$\vec{A}[\vec{r}(t), \Omega] = -\vec{\nabla}\phi. \quad (69)$$

From dimensional considerations we find that

$$\vec{A}[\vec{r}(t), \Omega] = \Omega^2 \vec{A}[\vec{r}(\Omega t), 1], \quad (70)$$

and that

$$\vec{A}[\lambda\vec{r}(t), \Omega] = \lambda\vec{A}[\vec{r}(t), \Omega], \quad (71)$$

Thus, $\vec{A}$ is necessarily homogeneous in $\vec{r}(t)$, and there is nothing to prevent us from having $\vec{A}$ that is linear in $\vec{r}(t)$. Theories with a linear kinetic term are rather more amenable to analysis; in particular their equation of motion is easy to derive from the action, when they are thus derivable; hence their heuristic

value. Henceforth we shall limit ourselves to such equations of motion, whose general form is

$$f(D/\Omega)D^2\vec{r}(t) = -\vec{\nabla}\phi. \quad (72)$$

In the present context the appropriate Newtonian limit is $\Omega \rightarrow 0$, if we want approximate Newtonian behavior for laboratory phenomena, where the typical frequencies would be much higher than Ω . Thus,

$$f(x) \xrightarrow{x \rightarrow \infty} 1. \quad (73)$$

Galilei invariance of $\vec{A}$ is assured if $f(x)$ is not singular at $x = 0$.

If we wish to follow this course beyond heuristics, and propound such theories as potential alternatives to dark matter, we have to note the following: For circular orbits in the mid-plane of a disk potential to be solutions of the equation of motion $f(x)$ must be an even function of x : $f(x) = h(-x^2)$. Then, eq. (72) gives for such orbits

$$h[(v/r\Omega)^2]v^2/r = \frac{d\phi}{dr}. \quad (74)$$

Flat rotation curves for disk galaxies result if

$$h(y) \approx y^{1/2}, \quad \text{for } y \ll 1. \quad (75)$$

The asymptotic rotational speed, V_∞ , is then related to the total galactic mass by $V_\infty^3 = MG\Omega$. This is at some odds with the observed Infra-red Tully-Fisher relation ($V_\infty^4 \propto L_{IR}$), if the IR luminosity of a galaxy is proportional to its mass (see point 2 in Sec. I). The value of Ω would have to be about $10^{-3}H_o$ to account for the mass discrepancy in galaxies (this value is approximately the orbital frequency of the sun in the Galaxy). Typical dynamical time scales in clusters of galaxies are much larger than in galaxies, so we would predict rather higher discrepancies in clusters. Not much of an Oort discrepancy is expected in the solar neighborhood, as the frequencies of the perpendicular motion are at least a few times higher than Ω , and due to the linearity of the kinetic term there is no direct effect of the low-frequency orbital motion on the perpendicular motion (there is only a shift in the equilibrium state for the oscillations). This linearity also assures the desired non-Newtonian center-of-mass motion of composite systems (stars, clouds, or atoms in galaxies, etc.) even though their components rattle inside them at high frequencies.

Phenomena in the laboratory or solar system are typified by frequencies, ω , much higher than Ω , and if the correction to Newtonian laws, in the limit $\omega \gg \Omega$, is of order $(\Omega/\omega)^2$ —as expected if $f(x)$ is even in its argument—it would not be detectable with present experimental capacity.

The evenness of f in its argument is tantamount to time-reversal (TR) symmetry of the equation of motion, and also to its derivability from a Lagrangian action with

$$L_k = \frac{1}{2}\vec{v}(t)h\left(\frac{\overleftrightarrow{DD}}{\Omega^2}\right)\vec{v}(t). \quad (76)$$

(Only the time-symmetric part of the Lagrangian affects the equation of motion, which is thus necessarily TR symmetric.)

Pais and Uhlenbeck[30] have discussed in detail the properties of a class of theories with Lagrangians equivalent to that of eq. (76), with $h(z)$ an entire (integral) function. They show that such theories suffer from the problems discussed in Sec. III. Our insistence on the particular Newtonian limit (73) (which they are not concerned with) puts us outside the province of their analysis: An entire function has an essential singularity at infinity (barring the conventional case of a constant f), and thus cannot have the Newtonian limit as required in eq. (73). (For a valid Newtonian limit, $f(z)$ must go to 1, as $z \rightarrow \infty$, in all directions in the complex plane, if the Lagrangian is to approach the Newtonian Lagrangian for all trajectories. For example, harmonic trajectories correspond to an imaginary argument of f ; time-exponential trajectories correspond to real ones, etc..) Our theory must then be non-local, indeed strongly non-local. In the present context this follows only from the required Newtonian limit; in the case of MOND, Galilei invariance, and the MOND limit were also brought to bear on the analogous conclusion. As we explain in Sec. III, the fact that our theory must not be entire in the time-derivative operator is a blessing: only such theories have a chance of being free of the difficulties that characterize higher-derivative theories.

VI. DISCUSSION

Examining modified laws of motion in the framework of the MOND scheme, we find that if the underlying kinetic action has the correct Newtonian limit, has the phenomenologically inspired MOND limit, and is Galilei invariant, the theory must be strongly non-local. It is this class of theories that we have concentrated on. They are not infinite-order limits of higher-derivative theories, which are known to have severe problems. We have demonstrated that MOND non-local theories may be formulated that are free of these problems. An attractive possibility is that MOND results as a nonrelativistic, small-scale expression of a fundamental theory by which inertia is a vestige of the interaction of a body with “the rest of the Universe”, in the spirit of Mach’s principle, or with an agent such as vacuum fields. The approximate equality of the acceleration “constant” of MOND, a_o , and cH_o , may be a witness. Such effective theories are usually strongly non-local.

We are not able yet to pinpoint a specific effective theory. The study of rotation curves—which provides the most detailed and clear-cut evidence for a mass discrepancy—is a strong test that is passed by all theories of the genre we consider here; but, it can only constrain the values of the kinetic action on circular orbits. Further constraints could come from the detection of mass discrepancies in light bending. Beyond these, existing observations of the mass discrepancy in galactic systems are hardly helpful in discerning a superior candidate theory. Additional discriminating power might come from the study of systems with non-circular motions, such as elliptical galaxies and

clusters of galaxies, because theories that predict the same circular motion usually disagree on non-circular trajectories.

Also potentially helpful is the study of a possible mass discrepancy manifested by the motion perpendicular to the plane in disk galaxies, which is a small perturbation on the circular motion. There may be such a mass discrepancy observed in the solar neighborhood, in the Milky Way (the Oort discrepancy), but the issue is still moot. Different theories of the present class predict different degrees of such a discrepancy. For example, it was pointed out[2] that eq. (3) predicts a discrepancy of magnitude $1/\mu(a_c/a_o)$ where a_c is the acceleration in the circular orbit (as long as the acceleration perpendicular to the disk is small compared with a_c). Analysis of small-perturbation, perpendicular motion in the theory defined by the action (40), gives a different value for the discrepancy: $1/[(1 + \hat{S}_k^c/2)F(S_k^c\omega^2/a_o^2)]$, where ω is the frequency of the perpendicular motion. Had ω been the frequency of the circular motion (actually $\ll \omega$), this would have given $1/\mu(a_c/a_o)$.

The examples we discuss are still formulated as test-particle dynamics. We cannot yet offer an example of a theory that fully accounts for the correct center-of-mass motion of composite objects: A composite body (e.g. a star), with constituents whose internal motion is Newtonian should still undergo a center-of-mass motion similar to that of a test particle, which may be well in the MOND regime. This we deem only a technical obstacle; a feeling that is supported by the following observation. We may formulate a MOND action as an integral over the Fourier transform, r_ω , of the trajectory, such that, in an extreme case, different frequencies are decoupled from each other, and for each frequency the motion depends on the value of $\omega^2 r_\omega/a_o$. Thus, the internal motions for which this value is large, obey approximate Newtonian dynamics, while the center-of-mass motion, characterized by a small value, is test-particle like. The theories discussed in Sec. V have this desired property. For various reasons we do not endorse such total spectral decoupling of the motion in the kinetic action (as occurs in the Newtonian action).

The action in eq. (40) does couple different frequencies, and also give a partial solution to the problem: Consider a trajectory of a constituent of a “star” that performs “galactic” motion with (say one) frequency ω_g and frequencies ω_i that describe the internal motion. We see from eq. (40) that, if the internal velocities are small enough compared with the galactic one, the value of S_k is determined by the “galactic” component, even if $\omega_i \gg \omega_g$ because F saturates at a value of 1 for large arguments. The condition for that is

$$(v_i/v_g)^2 \ll S_k \omega_g^2 / a_o^2. \quad (77)$$

Then, the “galactic” motion will be test-particle like, and the internal motions will depend on the value of $(S_k/v_i^2)(a_i^2/a_o^2)$, where $a_i \equiv v_i \omega_i$. This theory does not provide a fully satisfactory solution because it does not cover the case with internal velocities that are not smaller than the external one (motions within the atoms or nuclei are also internal). It does indicate, however that

the solution is within reach.

Another desideratum we have yet to implement is an extension of the theory to the relativistic regime: a matter-of-principle must, that is also needed to treat light bending. We feel that understanding the underpinning of MOND is a prior goal, and its achievement will surely bring about the understanding of relativistic MOND. We note in this connection that for a phenomenon to be both relativistic, and to involve accelerations of order a_o or below, it must also involve a length scale of cosmological magnitude. For instance, for the acceleration near the Schwarzschild radius of a mass M to be at a_o or below we must have this radius no smaller than the Hubble radius. Similarly, light trajectories in regions where the Newtonian acceleration is of order a_o , has a radius of curvature of the order of the Hubble radius. Thus, the relativistic-MOND regime falls in the large-scales regime. In light of the surmised cosmological origin of MOND this means that a relativistic extension of MOND might not simply hinge on the single acceleration constant a_o , which may, in fact, lose its role altogether.

Like all effective theories, ours must have validity bounds that stem from the approximations underlying the theory. These delimiters are usually not recognized from within the theory itself; but, sometimes, inconsistencies or other problems are vestiges of these approximations. Consider, for instance trajectories with constant velocity; in all the model actions we have studied they correspond to the extreme MOND limit in which the action vanishes. Clearly they do not have a Newtonian limit, which is one of the axioms of our treatment; we see a hint of a delimiter here. We should also not be dismayed if our effective (relativistic) theory fails to account for cosmology; as explained in Sec. I we certainly expect a delimiter there.

Finally, we list some differences between modifying gravity, and modifying inertia, beyond the fact that the latter pertains to arbitrary combinations of forces. We saw that conserved quantities and adiabatic invariants are different in the two approaches. In gravitational MOND theories, any isolated mass produces a confining gravitational potential: To obtain asymptotically flat rotation curves we must have an asymptotically logarithmic potential. It is not clear to us whether there is a general answer to the analogue question in our class of theories (i.e., are there unbound trajectories about an isolated mass?). The approximation underlying our theory may, in fact, break down for such unbound orbit, as indicated above.

APPENDIX A

We derive the general form of a Galilei-invariant, local, kinetic Lagrangian, L_k [which can be written as a function of $\vec{r}(t)$ and a finite number of its derivatives]. We make use of the commutation rules

$$\frac{\partial}{\partial \vec{r}} \frac{d}{dt} Q = \frac{d}{dt} \frac{\partial}{\partial \vec{r}} Q, \quad (A-1)$$

and

$$\frac{\partial}{\partial \vec{r}^{(i)}} \frac{d}{dt} Q = \frac{d}{dt} \frac{\partial}{\partial \vec{r}^{(i)}} Q + \frac{\partial Q}{\partial \vec{r}^{(i-1)}}, \quad i > 0. \quad (A-2)$$

[These stem from $\dot{Q} = (\vec{v} \cdot \frac{\partial}{\partial \vec{r}} + \vec{a} \cdot \frac{\partial}{\partial \vec{v}} + \dots)Q$.] For the equation of motion to be invariant under translations it is necessary (and sufficient) that $\vec{\nabla}_r L_k$ is some time derivative $\frac{d\vec{Q}}{dt}$. Taking the r -curl of this equality we get, with the aid of (A-1), that $\frac{d}{dt}[\vec{\nabla}_r \times \vec{Q}] = 0$. Now, when a quantity F depends on a finite number of derivatives of $\vec{r}(t)$, and $\frac{d}{dt}F = 0$ (i.e. F is constant along any trajectory), then F must be an absolute constant, independent of any of the variables: If F depends on $\vec{r}^{(0)}, \vec{r}^{(1)}, \dots, \vec{r}^{(k)}$ only, and $\frac{d}{dt}F = \vec{r}^{(1)} \cdot \frac{\partial F}{\partial \vec{r}^{(0)}} + \dots + \vec{r}^{(k+1)} \cdot \frac{\partial F}{\partial \vec{r}^{(k)}} = 0$, for any choice of $\vec{r}^{(0)}, \dots, \vec{r}^{(k+1)}$, then, clearly, all the partial derivatives of F must vanish. (Seen differently by noting that any two *finite* sets of derivatives are connected by some trajectory, so F must take the same value on any two sets.) Thus, $\vec{\nabla}_r \times \vec{Q}$ is a constant of the theory, and hence must actually vanish, because of the isotropy of the theory, which forbids the construction of a non-vanishing, constant vector. Thus, $\vec{Q} = \vec{\nabla}_r q$ for some q , and hence $\vec{\nabla}_r(L_k - \dot{q}) = 0$, and L_k is independent of $\vec{r}$ up to the time derivative $\dot{q}$, which we can discard.

Take then L_k that depends only on the derivatives of $r(t)$. For the equation of motion to be Galilei invariant we must have $\vec{\nabla}_v L_k = \frac{d\vec{Q}}{dt}$ for some $\vec{Q}$. The left-hand side does not depend on the variable $\vec{r}$, so take the derivative with respect to $\vec{r}$ to obtain $\frac{d}{dt}[\vec{\nabla}_r \otimes \vec{Q}] = 0$. As above, the tensor in parentheses must be a constant, and is dimensionless, so it must be $\alpha \delta_{ij}$, with α a dimensionless constant. Thus, $\vec{Q} = \alpha \vec{r} + \vec{U}$, where $\vec{U}$ is independent of $\vec{r}$. We then have $\vec{\nabla}_v L_k = \alpha \vec{v} + \frac{d\vec{U}}{dt}$. Taking the v -curl (which now commutes with the t -derivative on $\vec{U}$) we find that $\vec{\nabla}_v \times \vec{U}$ is a constant, and hence must vanish, and hence $\vec{U}$ is a v -divergence of some u that does not depend on $\vec{r}$. From all this we get

$$\vec{\nabla}_v(L_k - \frac{1}{2}\alpha v^2 - \dot{u}) = 0. \quad (A-3)$$

The expression in parentheses is, thus, some $\tilde{L}_k$ that does not depend on $\vec{r}$ or $\vec{v}$ as stated in eq. (33).

If L_k is not of finite order, but is a limit, for $N \rightarrow \infty$, of a sequence of Lagrangians, L_k^N , of finite order N , then, each of the L_k^N 's is of the form (33), and, hence, so is L_k (if the L_k^N 's are not themselves Galilei invariant, we can take their Galilei-symmetrized parts, and they too go to L_k for $N \rightarrow \infty$).

APPENDIX B

Here we derive expression (29) for the energy function, which applies whenever this function can be defined. Starting with the general expression for the energy, eq. (28), and shifting time derivatives from the second factor

to the first, we can write (for $m \geq 2$)

$$\vec{r}^{(i-m+1)} \cdot \left[\frac{\partial L}{\partial \vec{r}^{(i)}} \right]^{(m-1)} = \dot{q}_i + (-1)^{m-1} \vec{r}^{(i)} \cdot \frac{\partial L}{\partial \vec{r}^{(i)}}, \quad (B-1)$$

with

$$q_i = - \sum_{l=1}^{m-1} (-1)^l \vec{r}^{(i-m+l)} \cdot \left[\frac{\partial L}{\partial \vec{r}^{(i)}} \right]^{(m-l-1)}. \quad (B-2)$$

Upon taking the double sum in eq (28), the first term in eq.(B-1) gives $\dot{Q}$ with Q given by eq. (30) (after some rearrangement and summation). The second term in eq.(B-1) gives $\sum_i i \vec{r}^{(i)} \cdot \frac{\partial L}{\partial \vec{r}^{(i)}}$, which we now evaluate. Imagine that L depends on a number of dimensional constants C_ℓ , $\ell = 1, \dots, I$, with dimensions $(length)^{\alpha_\ell} \times (time)^{\beta_\ell}$, respectively. The Lagrangian itself has dimensions $(length)^2 \times (time)^{-2}$.

Considering the redefinition of the units of time we have

$$L[\lambda^{\beta_1} C_1, \dots, \lambda^{\beta_I} C_I, \lambda^{-1} \vec{r}^{(1)}, \dots, \lambda^{-i} \vec{r}^{(i)}, \dots] = \lambda^{-2} L[C_1, \dots, C_I, \vec{r}^{(1)}, \dots, \vec{r}^{(i)}, \dots]. \quad (B-3)$$

Taking the derivative with respect to λ at $\lambda = 1$, we find, as in the Euler theorem for homogeneous functions,

$$\sum_{\ell=1}^I \beta_\ell C_\ell \frac{\partial L}{\partial C_\ell} - \sum_{i=1}^N i \vec{r}^{(i)} \cdot \frac{\partial L}{\partial \vec{r}^{(i)}} = -2L. \quad (B-4)$$

Putting eq.(B-4) together with the above, we get

$$H = L + \sum_{\ell=1}^I \beta_\ell C_\ell \frac{\partial L}{\partial C_\ell} + \dot{Q}. \quad (B-5)$$

We separate the Lagrangian into kinetic and potential parts $L = L_k - \phi$; ϕ , which is taken to depend only on $\vec{r}$, must be of the form $C f(\vec{r}/r_0)$, where C has dimensions of velocity squared, and r_0 those of length. Thus the contribution of ϕ to the sum in (B-5) is 2ϕ , and none to Q . We can then also write

$$H = L_k + \sum_{\ell=1}^I \beta_\ell C_\ell \frac{\partial L_k}{\partial C_\ell} + \phi + \dot{Q}, \quad (B-6)$$

and specializing to the case of a single constant a_o ($\beta = -2$), we get eq. (29).

Another useful relation is obtained by considering, instead, redefinition of the units of length, which gives

$$L_k[\lambda^{\alpha_1} C_1, \dots, \lambda^{\alpha_I} C_I, \lambda \vec{r}^{(1)}, \dots, \lambda \vec{r}^{(i)}, \dots] = \lambda^2 L_k[C_1, \dots, C_I, \vec{r}^{(1)}, \dots, \vec{r}^{(i)}, \dots], \quad (B-7)$$

from which follows, as above,

$$\sum_{\ell=1}^I \alpha_\ell C_\ell \frac{\partial L_k}{\partial C_\ell} + \sum_{i=1}^N \vec{r}^{(i)} \cdot \frac{\partial L_k}{\partial \vec{r}^{(i)}} = 2L_k. \quad (B-8)$$

References

1. M. Milgrom, *Astrophys. J.* **270**, 365 (1983).
2. M. Milgrom, *Astrophys. J.* **270**, 371 (1983).
3. M. Milgrom, *Astrophys. J.* **270**, 384 (1983).
4. S.M. Kent, *Astronom. J.* **93**, 816 (1987).
5. M. Milgrom, *Astrophys. J.* **333**, 689 (1988).
6. R.H. Sanders, *Astron. Astrophys. Rev.* **2**, 1 (1990).
7. K.G. Begemen, A.H. Broeils, and R.H. Sanders, *Mon. Not. R. astr. Soc.* **249**, 523 (1991).
8. G. Lake, *Astrophys. J.* **345**, L17 (1989).
9. M. Milgrom, *Astrophys. J.* **367**, 490 (1991).
10. R.B. Tully and J.R. Fisher, *Astron. Astrophys.* **54**, 661 (1977). M. Aaronson and J. Mould, *Astrophys. J.* **265**,1 (1983). M. Aaronson, G. Bothun, J. Mould, J. Huchra, R.A. Schommer, and N.E. Cornell, *Astrophys. J.* **302**, 536 (1986).
11. M. Milgrom, *Astrophys. J.* **287**, 571 (1984).
12. S.M. Faber and R.E. Jackson, *Astrophys. J.* **204**, 668 (1976).
13. C. Carignan, R. Sancisi, and T.S. van Albada, *Astron. J.* **95**, 37 (1988). C. Carignan and S. Beaulieu, *Astrophys. J.* **347**, 192 (1989). G. Lake, R.A. Schommer, and J.H. van Gorkom, *Astron. J.* **99**, 547 (1990). D. Puche, C. Carignan, and A. Bosma, *Astron. J.* **100**, 1468 (1990). M. Jobin and C. Carignan, *Astron. J.* **100**, 648 (1990).
14. C. Pryor and J. Kormendy, *Astron. J.* **100**, 127 (1990). M. Mateo, E. Olszewski, D.L. Welch, P. Fischer, and W. Kunkel, *Astron. J.* **102**, 914 (1991).
15. R.A. Fish, *Astrophys. J.* **139**, 284 (1964).
16. P.C. van der Kruit, *Astron. Astrophys.* **173**, 59 (1987).
17. M. Milgrom, *Astrophys. J.* **338**, 121 (1989). D.M. Christodoulou, *Astrophys. J.* **372**, 471 (1991).
18. See e.g. J.A. Tyson, F. Valdes, and R.A. Wenk, *Astrophys. J. Lett.* **349**, L1 (1990). J.A. Tyson, *Phys. Today*, June, 24 (1992).
19. J.D. Bekenstein and M. Milgrom, *Astrophys. J.* **286**, 7 (1984).

20. J.D. Bekenstein, in *General Relativity and Astrophysics*, eds. A. Coley, C. Dyer, and T. Tupper, (World Scientific, Singapore, p. 64) (1988). J.D. Bekenstein, Phys. Lett. B, **202**, 497 (1988). R.H. Sanders, Mon. Not. R. astr. Soc. **223**, 539 (1986). R.H. Sanders, Mon. Not. R. astr. Soc. **235**, 105 (1988). J.D. Bekenstein, Sixth Marcel Grossman Conference on General Relativity, Kyoto (1991).
21. M. Milgrom, Comm. Astrophys. **13**, 215 (1989).
22. J.P. Huchra, Science **256**, 321 (1992).
23. S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989).
24. S.M. Carroll, W.H. Press, and E.L. Turner, Ann. Rev. Astron. Astrophys. **30**, 499 (1992).
25. See e.g. P.J.E. Peebles, Astrophys. J. **284**, 439 (1984); G. Efstathiou, W.J. Sutherland, and S.J. Maddox, Nature **348**, 705 (1990).
26. See, e.g. W. Misner, K.S. Thorne, and J.A. Wheeler *Gravitation*, W.H. Freeman, N.Y. (1973).
27. J.A. Wheeler and R.P. Feynman, Rev. Mod. Phys. **21**, 425 (1949).
28. See e.g. A.O. Caldeira and A.J. Legget, Ann. Phys. **149**, 347 (1983) and references therein.
29. M. Ostrogradski, Mem. Act. St. Petersburg **VI4**, 385 (1850).
30. A. Pais and G.E. Uhlenbeck, Phys. Rev. **79**, 145 (1950).
31. X. Jaén, J. Llosa, and A. Molina, Phys. Rev. **D 34**, 2302 (1986).
32. D.A. Eliezer and R.P. Woodard, Nucl. Phys. **B325**, 389 (1989).
33. J.Z. Simon, Phys. Rev. **D 41**, 3720 (1990).